

**LIST OF MEDICAL PROBLEMS, THEIR
PATHOGENIC FACTORS AND THEIR TREATMENT
USING “SELECTIVE DRUG UPTAKE
ENHANCEMENT” (SDUE) METHOD**

©Copyright Yoshiaki Omura M.D., Sc.D. 1998-2006

Presented by Yoshiaki Omura M.D., Sc.D. of New York

During His Seminars and Workshops

SYMPTOMS OR DISEASE	MOST COMMON CAUSATIVE AGENT Indicated by Bi-DORT Procedure <i>※太字のものは RCS があるもの</i>	MEDICATIONS All medications are 4 times per day with SDUE unless otherwise noted	AREA OF SELECTIVE DRUG UPTAKE STIMULATION
Abortion (spontaneous) 流産(自然流産)	Cytomegalovirus Infection Chlamydia trachomatis (one major cause) Hg	EPA/DHA Doxycycline or Azithromycin Cilantro	Appropriate area on hand chart
Achalasia (アカラシア) (esophageal stricture @ sphincter) (食道狭窄@. 括約筋)	CMV Cheese intake	EPA/DHA	Third segment of middle finger in midline
ALS (Amyotrophic Lateral Sclerosis) (Lou Gherig Disease) ALS (筋萎縮[性]側索硬 化[症]) (レー・ゲーリグ病)	Localized Deposits of Hg , and Chlamydia trachomatis , Cytomegalovirus in parietooccipital (頭頂後頭 の) area of brain genitourinary (尿生殖器 の) area; Candida Albicans	Cilantro, Doxycycline EPA/DHA	Affected areas on hand chart
Allergies (アレルギー)	Chlamydia trachomatis infection in mucous membranes of sinuses, nose, mouth, throat, etc	Doxycycline or Azithromycin	Sinus/Nose/ Mouth/Throat (first segment of Middle finger- Ventral(腹面)); Lung areas of hand chart(third segment-sides)
Alzheimer's Disease Pre-Alzheimer's Disease (アルツハイマー病) (アルツハイマー前状態)	Increased Al, Hg & Pb Amyloid-β(1-42) and Decreased Acetylcholine L-Glutamic Acid Mg in Hippocampus. DHEA Cytomegalovirus (most common virus found) Tau Protein HHV-6 &/or7 HSV-I Chlamydia trachomatis Mycobacterium tuberculosis	Cilantro EPA/DHA Folic Acid α-Lipoic Acid Acetyl-L-Carnitine DHEA	First segment of middle finger (brain)

	L-homocystine 8-OH-dG		
Asthma (喘息)	Chlamydia trachomatis Chlamydia Pneumonia α-Streptococcus Coxsackie virus etc. Identify specific organism/s and prescribe accordingly	Doxycycline (100 mg bid for 2-3 mos) Trimox EPA/DHA	Lower 1/2 of first segment of middle finger, second & third segment of middle finger (all surfaces)
Atrial Fibrillation, Auricular Fibrillation (心房[性]細動)	Chlamydia trachomatis or Borrelia Burgdorferi in SA node of heart	Doxycycline Trimox	Third segment of middle finger (ventral surface)
Autism (自閉症) (Childhood Alzheimers)	Increased Al, Pb, Hg Decreased Acetylcholine CMV 8-OH-dG HHV-6 & HHV-7	Cilantro EPA/DHA Folic Acid	First segment middle finger (all surfaces)
Bad Breath (combined with Gastric problems) (口臭)	Helicobacter Pylori	Trimox	Mouth and gastric area of hand chart
Bell Palsy(ベル麻痺) Peripheral Facial Paralysis (末梢性顔面麻痺)	HSV-I (major cause) Borrelia Burgdorferi Al, Pb, Hg	EPA/DHA Trimox Cilantro	First Segment of Middle Finger (ventral surface)
Black Hairy Tongue 黒[色]舌, 黑毛舌	Fungus	Diflucan 100 mg q.d. for 2+ weeks	Tongue(also use tongue scraper)
Blood Pressure 血压	Infection of cardiovascular representation area of Medulla oblongata & heart w/ HSV-I &/or HSV-II ; or Chlamydia trachomatis in Aortic arch. CMV; Epstein-Barr Virus	EPA/DHA Cilantro if heavy metals are present Doxycycline	Back of First Segment of Middle Finger below nail. Ventral surface of 3rd segment of middle finger
Bulimia 過食症	Heavy metals in brain	EPA/DHA & Cilantro	First segment of middle finger
Cancer (癌)	Increased Hg, Integrin $\alpha 5\beta 1$ & Oncogene C-fos Ab2 ; Less than 1 pg of Acetylcholine 8-OH-dG Presence of a virus DHEA Increased Telomere (TTAGGG) in cancer cell	EPA/DHA & Cilantro Folic Acid α -Lipoic Acid Acetyl-L-Carnitine DHEA	Organ representation area of affected organ
Cancer - Cervical (子宫癌)	Human Papilloma virus HPV16 or HPV18 8-OH-dG DHEA Telomere	EPA/DHA & Cilantro Folic Acid DHEA	Appropriate area on hand chart
Cancer - Esophageal (食道癌)	CMV 8-OH-dG Cheese intake DHEA Helicobacter Pylori Telomere	EPA/DHA & Cilantro Folic Acid DHEA	Second & third segment of middle finger in midline
Cancer - Gastric	CMV 8-OH-dG DHEA	EPA/DHA &	Appropriate organ

(胃癌)	Cheese intake <i>Helicobacter Pylori</i>	Salt intake <i>Mycobacterium</i> <i>Tuberculosis</i> >1µg <i>Mycobacterium Avium</i> >1µg	Cilantro Folic Acid DHEA Trimox	representation area on hand chart
Chronic Cough (慢性的な咳)	<i>Mycobacterium</i> <i>Tuberculosis</i> >1µg <i>Mycobacterium Avium</i> >1µg	Isoniazid 300 mg bid & Refampin 150 mg bid (human) Isoniazid + B6(Avium)	Third segment of middle finger (sides)	
Chronic Fatigue Syndrome (慢性疲労症候群)	Decreased circulation to the Pons causing decreased Serotonin and Nor-epinephrine ; Epstein Barr Virus ; CMV , HHV-6 & 7 , AIDS (HIV) <i>Mycoplasma pneumoniae</i>	Massage vertebral artery@C5; EPA/DHA, Cilantro if metal deposit	First Segment of Middle Finger (dorsal surface)	
Circulation (Decreased) 循環障害	Electromagnetic Fields; Increased Thromboxane B2 ; Any Infection (virus, bacterial)	Laser to St. 36, medulla oblongata and dorsal surface of first phalanx of middle finger	Acupuncture to area of decreased circulation every 2-3 days	
Convulsions 痙攣 (顔, 体幹, 四肢の激しい痙攣または連続的痙攣)	CMV in brain HHV-6 in brain	EPA/DHA & Cilantro	First segment of middle finger	
Crohn Disease Regional Enteritis クローン病、限局性腸炎	<i>Chlamydia trachomatis</i> Virus infection; TB infection	Doxycycline EPA/DHA, Cilantro if metal deposit	Colon area on hand chart	
Decreased Acetylcholine in Brain 脳内アセチルコリンの減少	Most Infections; Electromagnetic fields Hg, Pb, Al	Appropriate antibiotic or EPA/DHA w/Cilantro	First Segment of Middle Finger (Dorsal surface lower half)	
Depression うつ病, 抑うつ[症]	Decreased Serotonin in Raphe Nuclei(縫線核) in the Pons due to Vasoconstriction (血管収縮) of vertebral A(椎弓)&/or Basilar A(基底角); or, a virus or bacterial infection in Pons & medulla (髓質) due to Metal deposits	First O-ring is weak EPA/DHA & Cilantro if viral Trimox or Doxycycline if bacterial	Massage vertebral Artery @C5 if due to vasoconstriction Appropriate meds if due to infection	
Diabetes 糖尿病	Infection in pancreas of Cytomegalovirus plus a bacteria; <i>Chlamydia trachomatis</i> Pseudomonas aeruginosa α-Streptococcus <i>Candida Albicans</i> in pancreas 歯周病菌 Decreased Insulin	EPA/DHA,Cilantro Doxycycline Trimox Trimox & Cilantro	Pancreas area on hand chart	

Diarrhea (Intractable) 下痢	Chlamydia trachomatis in descending colon; Tuberculosis(human)	Doxycycline (100 mg qid for 2-3 mos)	Colon area on Hand chart
Dementia 痴呆	Decrease Acetylcholine Aluminum in brain β-Amyloid(1-42) Tau	Cilantro	First segment of middle finger
Diverticulitis 憩室炎	Virus Infection in colon	EPA/DHA, Cilantro	Appropriate organ representation area
Dysphagia 嚥下困難, 嚥下障害	Cytomegalovirus in Esophagus Coxsackie's virus (occasionally)	EPA/DHA & Cilantro	Midline of middle segment of finger (ventral surface)
Dystonia (Vallismus) ジストニア	Viral Infection in brain HSV I, II;HHV-6, 7 Cytomegalovirus , &/or Chlamydia trachomatis or other bacteria Decreased GABA (gamma aminobutyric acid) in Cerebellum and decreased Acethylcholine & Dopamine ; Increased metal deposits(Al,Hg, Pb)	EPA/DHA, Cilantro Appropriate Antibiotic/s	First segment of middle finger (all surfaces)
Electromagnetic Field Hypersensitivity 電磁波過敏症候群	High concentration of heavy metals(Hg, Pb) & aluminum(Al) in Hippocampus area of brain; Decreased Acetylcholine	EPA/DHA, Cilantro	First segment of middle finger (all surfaces)
Endometriosis 子宮内膜症	HHV-6 in uterus Multiple types of Virus	EPA/DHA	Appropriate Area on hand chart
Environmental Hypersensitivity (chemicals, dust, pollen, electricity) 化学物質過敏症 (化学物質、ほこり、花粉、電気)	Chlamydia trachomatis infecting mucosa of mouth, nose and throat, High concentrations of metals(Al, Hg, Pb)	Doxycycline Azithromycin (Zythromax) EPA/DHA & Cilantro	Appropriate organ representation area on hand chart
Epstein-Barr	Epstein-Barr virus	EPA/DHA	
Esophageal Reflux (precursor to esophageal Cancer) 食道逆流 (食道癌の前駆 症状)	Cytomegalovirus in esophagus Excessive intake of munster cheese Excessive intake of nuts	EPA/DHA, Cilantro	Midline of 2nd &3rd segments of middle finger (ventral surface)
Face Muscle Spasm (tics) 顔面痙攣, 顔面筋痙攣	Borrelia Bufgdofieri (Lyme's Disease)	EPA/DHA Timox	First segment of middle finger (ventral surface)

Facial Paralysis 顔面神経麻痺	Virus	EPA/DHA	First segment of middle finger (ventral surface)
Fibromyalgia 線維筋痛症 (筋骨格起源であるが原因不明の慢性痛を呈する症候群)米国リウマチ協会は診断基準によれば、その疼痛は体軸部位(頸、胸、腰椎または前胸部)とともに体の両側全身にみられ、さらに18か所の特定の圧痛点のうち少なくとも11か所に圧痛を認めることが必要である	Decreased and increased Nor-epinephrine Thromboxane B2 in Pons; Mixed viral & bacterial infection; HSV-I & II; Chlamydia trachomatis, Borrelia Burgdorferi (Lyme's Disease)	EPA/DHA, Cilantro Doxycycline Trimox 500 mg	Area of greatest symptomatology
Frequent Sneezing 頻繁なクシャミ	Chlamydia trachomatis	Doxycycline	First segment of middle finger (ventral surface)
Fungus Infection 菌感染症	Multiple types of fungi	Diflucan	Appropriate area of hand chart
Gillian Barre Syndrome ギランバレー症候群	Virus	EPA/DHA, Cilantro	First segment of middle finger (ventral surface)
Glaucoma (Trachoma) 緑内障(トラコーマ) 網膜色素変性症	Chlamydia trachomatis in retina	Doxycycline	First segment of middle finger (ventral surface)
Headaches (bilateral) 頭痛(両側[性]の)	HSV-I on one side and HSV-II on other side of brain; Human TB in brain; Chlamydia Trachomatis in brain	EPA/DHA & Cilantro Isoniazide with or w.o. Refampin; Doxycycline or Azithromycin	First segment of middle finger
Headaches(unilateral) 頭痛(片側[性]の、一側[性]の)	HSV-I	Cilantro EPA/DHA	Appropriate Area on hand chart
Hearing Loss 難聴	Pseudomonas HSV-I or HSV-2	Trimox	First segment of middle finger (sides)
Heart Problems Cardiovascular Disease (arrhythmias, etc) 心臓病(不整脈等)	Chlamydia Pneumoniae Chlamydia trachomatis virus & Bacterial Infection Coxsackie's virus PLGF HHV-6 CRP Lp(a) PAI-1 L-homocystine 8-OH-dG 歯周病菌 DHEA	B12 & Folic Acid Doxycycline EPA/DHA & Cilantro	Third segment Of middle finger (ventral surface) Acupuncture to St36
Hepatitis B-type C-type	Multiple type of virus Anti-HBs Anti-HBe Anti-HCV recombinant HBe	EPA/DHA	Liver area on hand chart
Herpetic Neuralgia (ヘルペス神経痛)	Virus Substance P Herpes Virus Series	EPA/DHA& Cilantro	Appropriate area on hand chart
Herpes Zoster (Shingles)	HSV-I HSV-II	Cilantro	Appropriate area

ヘルペスゾスター	Varicella Zoster Virus	EPA/DHA	on hand chart
High Blood Pressure (See Blood Pressure) 高血圧	Chlamydia trachomatis in aortic arch. Infection in medulla oblongata	Doxycycline EPA/DHA& Cilantro	Third segment of middle finger (ventral surface) & first segment of middle finger below fingernail (dorsal surface)
Hydrocephalus 水頭[症]	Toxoplasma Gondii	Bactrium DS	First segment of middle finger (dorsal surface)
Hypersensitivity to pain 疼痛過敏症	Decreased Serotonin in Raphe Nuclei of the Pons due to bacteria and virus infection; vasoconstriction of the Vertebral or Basilar A; or, to injury to Vertebral A, due to chiropractic treatment	Massage Vertebral artery (椎骨動脈) @C5; Appropriate antibiotic/s; EPA/DHA with Cilantro if indicated	First segment of middle finger (dorsal surface)
Hypersensitivity to: Pollen、Dust、Chemicals 花粉、ホコリ、化学物質に対する過敏症	Chlamydia trachomatis Histamine Formaldehyde Bisphenol A etc.	Doxycycline	Appropriate area of hand chart
Hypersensitivity (increased sensitivity to touch)接触性過敏症	HSV-I or HSV-II Heavy Metals	EPA/DHA Cilantro	Appropriate area of hand chart
Infected Joints 関節炎 (See Rheumatoid A) 慢性関節リウマチ参照	Mixed infection of Borrelia Burgdorferi (Lyme Disease) and Chlamydia trachomatis	Trimox Doxycycline	Appropriate joint area of hand chart
Infertility 不妊症	Chlamydia trachomatis 歯周病菌	Doxycycline	Appropriate area of hand chart
Insomnia 不眠[症]	Decreased Serotonin in brain and Raphe Nuclei of the Pons due to bacteria and virus infection; check all areas of the brain; vasoconstriction of the Vertebral or Basilar A; or, injury to Vertebral A, due to chiropractic treatment	Massage vertebral artery@C5 Appropriate antibiotics and/or EPA/DHA with Cilantro if indicated	First segment of middle finger (all surfaces)
Intractable Face or Tooth Pain 難治性顔面痛 or 歯痛	HSV-I and HSV-II	EPA/DHA & Cilantro	First segment of middle finger (ventral surface)
Infectious Mononucleosis 感染性单核細胞症 (循環血液中に、单核白血球が異常に多数存在する疾患)	Epstein Barr Virus	EPA/DHA & Cilantro	

Joint Pain - Multiple Joints (See Infected Joints) 関節痛	Borrelia Burgdorferi Chlamydia trachomatis Metal Deposits Uric Acid	Trimox, Doxycycline (for 2-3 mos);Cilantro	Affected Joints on hand chart
Lupus 狼瘡	Viral infection, Hg, Al, Pb deposits	EPA/DHA Cilantro	Affected area on hand chart
Lyme's Disease (ライム病)	Borrelia Burgdorferi	Trimox (500mg qid) or Tetracycline (500mgqid)(LederleCo.)	Appropriate Joint area on hand chart
Macular Degeneration 黄斑変性 (後極部を主に侵す眼の変性. 最も多い加齢性黄斑変性)	Chlamydia trachomatis Chlamydia Pneumoniae Herpes, Other viruses	EPA/DHA Cilantro	First segment of Middle finger (ventral surface)
Maxillary Sinus Pain 上顎洞痛	HSV-I	EPA/DHA & Cilantro	First segment of middle finger on same side. (ventral surface)
Memory Loss — Short Term 記憶障害	Large deposits of Hg, Pb & Al in Hippocampus of Brain; Decreased Acetylcholine (<500 µg) in brain; frequent viral or bacterial infection (CMV; HHV 6, 7;TB)	Cilantro	First segment of middle finger on side closest to thumb
Mental Development retarded 知能発達障害	Same Memory Loss Decreased Acethylcholine in Brain; HSV-I & II in brain & Spinal Cord; Coxsackie's virus ; Toxoplasmosis in newborn	EPA/DHA & Cilantro	First segment of middle finger for the Brain (all surfaces); Midline of middle finger for spinal cord (dorsal surface)
Microcephalus	Toxoplasma Gondii in brain	Bactrim DS	First segment of middle finger
Migraines with Aura 前兆のある偏頭痛	Infection with vascular involvement		
Migraines without Aura 前兆なしの偏頭痛	Infection Scalp Muscles		
Mitral Valve Prolapse 僧帽弁逸脱[症]	Chlamydia Trachomatis or Virus infection	Doxycycline EPA/DHA	Third segment of middle finger (ventral surface)
Movement Disorders 運動障害	Cytomegalovirus in brain	Cilantro with EPA/DHA	First segment of middle finger
Multiple Sclerosis 多発[性]硬化症	Chlamydia trachomatis with HSV-I or Mumps virus in sensory motor cortex of brain or in spinal cord and Genitourinary area. Hg Deposits	Trimox Doxycycline Cilantro EPA/DHA	First segment of middle finger for the Brain (all surfaces); Midline of middle finger for spinal cord

	Cytomegalovirus Borrelia Burgdorferi Candida Albicans		(dorsal surface)
Myasthenia Gravis 重症筋無力症	Cytomegalovirus Hg &/or Pb	Cilantro	First segment of middle finger (all surfaces)
Obsessive Compulsive Disorders — Bulimia 強迫観念にとらわれた異常過食症	Heavy Metals & Al in brain; viral or bacterial infection in brain	Cilantro	First segment of third finger
Osteoarthritis 変形性関節症	Chlamydia trachomatis Borrelia Burgdorferi Uric acid	Doxycycline Trimox or Tetracycline	Appropriate area on hand chart
Pain on one side of body 片側面の疼痛	HSV-I or II (occasionally)	EPA/DHA & Cilantro	Appropriate area on hand chart
Pancreatic Infection 膵臓の感染症	Cytomegalovirus Chlamydia trachomatis Pseudomonas 歯周病菌	EPA/DHA	Appropriate area on hand chart
Paralysis of Skeletal Muscles 骨格筋の麻痺	Cytomegalovirus (most common) Decreased Acetylcholine Decreased Dopamine Increased GABA (gamma aminobutyric acid) HSV-I, Hg, Pb deposits	EPA/DHA Cilantro Electric Stimulation	Appropriate area(brain area if due to stroke)
Parkinson's Disease (see Tremor) パーキンソン病 (振せん, 震え 参照)	Cytomegalovirus and Metal deposits in Caudate Nucleus(尾状核)	Cilantro EPA/DHA	First segment of middle finger at side of nail, nail, and the area below nail to first joint
Pemphigus 天疱瘡	<u>Bacterial infection</u>	Bactrim DS Doxycycline (Do not use together)	Appropriate area on hand chart
Post Herpetic Neuralgia ヘルペス後神経痛	HSV-I & II; Metal Deposits in affected area	EPA/DHA, Cilantro	Appropriate area on hand chart
Premenstrual Pain 月経前痛	Chlamydia trachomatis	Doxycycline	Appropriate area on hand chart
Pseudomonas Infection 緑膿菌感染	Pseudomonas	Trimox	Appropriate area on hand chart
Psoriasis 乾癬	<u>virus and/or bacterial infection;</u> Hg, Pb, Al deposits	EPA/DHA; Trimox Cilantro	Appropriate area on hand chart
Reynaud's Disease	Chlamydia trachomatis	Doxycycline	Most affected

レイノー病	or HHV-6 or HHV-7 in circulatory system; HSV-I & II; Helicobacter pylori in gastrointestinal system	EPA/DHA Cilantro Trimox	area on hand chart
Rheumatoid Arthritis 慢性関節リウマチ	A mixed infection of Chlamydia trachomatis and Borrelia Burgdorferi (Lyme's Disease) uric acid	Trimox DHA/EPA, Cilantro Doxycycline	Most affected area on hand chart
Scleroderma 強皮症, 硬皮症	Hg, Al deposits	EPA/DHA, Cilantro	Appropriate area on hand chart
Seizures (convulsions) 脳卒中 (ケイソ、ヒキツク)	HHV-6 in brain	EPA/DHA & Cilantro	First segment of middle finger
Shingles (see Herpes Zoster) 带状疱疹 (ヘルペスゾスター参照)	HSV-I or II; Herpes Zoster	EPA/DHA & Cilantro	Appropriate area on hand chart
Sinus Allergy 洞アレルギー	Chlamydia trachomatis		First segment of middle finger (ventral surface)
Sinus Pain 洞の痛み	HSV-I if on one side; HSV-I & II if on both sides	EPA/DHA, Cilantro	First segment of middle finger (ventral surface)
Sjögren's Disease シエーゲン症候群	Cytomegalovirus in Cerebellum, Pons and Medulla. Hg, Al	EPA/DHA, Cilantro	Most symptomatic area on hand chart
Sneezing クシャミ	Chlamydia Trachomatis in nose and throat	Doxycycline	First segment of middle finger (ventral surface)
Spasm of Diaphragm 横隔膜のけいれん	Chlamydia Trachomatis	Doxycycline (for 2-3 mos)	Appropriate area on hand chart
Spasm of Intercostal Membranes 肋間膜のけいれん	HSV-I or II Chlamydia trachomatis	EPA/DHA with Cilantro Doxycycline for 2,3 mos Without Cilantro	Appropriate area on hand chart
Stomach Infection 胃の感染	Helicobacter Pylori	Trimox	Appropriate area on hand chart
Stroke 脳卒中	Chlamydia Pneumoniae Decreased Acetylcholine in brain; HSV-I in motor cortex (運動皮質); Pb in motor cortex; Cytomegalovirus ;	Trimox Cilantro EPA/DHA	First segment of middle finger. (all surfaces) Acupuncture lo St.36, GB20, GB21 & dorsal Surface of cervical spine.

Thyroid Problems 甲状腺障害	Viral Infection/s TSH T3 T4 Parathyroid hormone	EPA/DHA with Cilantro	Second segment of middle finger
Tinnitus 耳鳴(じめい), 響鳴	Multiple causes: -Pseudomonas -HHV-6 or 7 -Chlamydia Trachomatis -Metal Deposits -TB in ear	Trimox EPA/DHA Doxycycline Cilantro Isoniazid and B-6	First segment of middle finger (sides)
Tooth Pain 歯痛	Alpha Streptococcus; Helicobacter pylori; Chlamydia Trachomatis; Borrelia Burgdorferi in periodontal membrane space	EPA/DHA with Cilantro Trimox, Bacterim DS Doxycycline without Cilantro	First segment of middle finger (sides and ventral surfaces)
Torticollis 痙性斜頸	HSV-I with or without HHV,6 &/or7	EPA/DHA with Cilantro Acupuncture to belly of SCM	Second segment of middle finger (sides)
Toxoplasmosis トキソプラズマ症	Toxoplasma Gondii is an Intracellular Parasite from Pork(20%), chicken and beef(5-10%)	Bacterim DS- 300 mg bid	Scan with X-Y axis laser to locate affected organ; appropriate area on hand chart
Trigeminal Neuralgia 三叉神経痛	Chlamydia Trachomatis HSV-I (most common) HSV-II	Trimox EPA/DHA with Cilantro	First segment o middle finger (same side as pain)
Tuberculosis (human) (constipation & diarrhea) ヒト型結核 (便秘&下痢)	Mycobacterium Tuberculosis	Isoniazid(300 mg bid for 2-3 months), Refampin (150 mg bid), B6(25 mg bid);Do not take EPA/DHA or Doxycycline	Third segment of middle finger (sides)
Tuberculosis(Avium) トリ型結核	Mycobacterium Avium	Isoniazid (3 00 mg bid for 2-3 months), &B6(25 mg bid); Do not take EPA/DHA or Doxycycline	Third segment of middle finger (sides)
Unilateral Face Pain (severe) 片側顔面痛	HSV-I(>25 ng); HSV-II(rarely); Metal Deposits	Cilantro	First segment of middle finger (same side as pain)

Uterine Cancer 子宮癌	Human Papilloma virus HPV16 or HPV18 DHEA 8-OH-dG Telomere	EPA/DHA with Cilantro	Appropriate area in hand chart
Vaginal Fungus Infections 膣真菌症	Yeast, antibiotics Metal deposits	Diflucan, 1 tab for 7 days (repeat if necessary) Cilantro	Appropriate area in hand chart
Varicose Veins 拡張蛇行靜脈	HHV-6 &/or 7; Chlamydia trachomatis	Doxycycline EPA/DHA, Cilantro	Appropriate area in hand chart
Vision Loss (Trachoma) 視覚障害 (トロコーマ)	Chlamydia trachomatis in retina Borrelia Burgdorferi Chalmydia Psittaci	Doxycycline(2-3 mos) Trimox	First segment of middle finger (ventral surface)
Vertigo めまい(眩暈)	Mycobacterium Tuberculosis(TB) in ear TB in cerebellum	Isonizid (300mg bid) with or w/o Refampin (150 mg bid); B6(25 mg bid)	First segment of middle finger