癌に関連したRCS

ORT生命科学研究所

Criteria used for the Bi-Digital O-Ring Test

Screening for Pre-Cancer & Cancer

- ◎① Oncogene C-fos Ab2 ↑ ↑ ↑ ↑ ↑
 - ◎② Integrin $\alpha_5\beta_1$ ↑ ↑ ↑ ↑ ↑
 - ◎③ Hg (Mercury) ↑ ↑ ↑ ↑ ↑
 - ◎④ Acetylcholine (Ach) $\doteq < 1/1000 \times \text{Normal}(\downarrow \downarrow \downarrow \downarrow \downarrow)$
 - ◎⑤ Viral Infection ↑ ↑ ↑ ↑ ↑
 - ⑥ NO (Nitrogen monoxide) $\doteq < 1/100 \times \text{Normal}(\downarrow \downarrow \downarrow \downarrow \downarrow)$
 - ⑦ D-Glucose ↑ ↑ (Max:2 × Blood Glucose Concentration)
 - ⑧ Rb (Ab-8) ↑ ↑ ↑ ↑ ↑
 - ⑨ p53(Ab-5) ↑ ↑ ↑ ↑ ↑
 - ⑩ Telomere-1(TTAGGG) ↑ ↑ (Max:5 × Normal Tissue Telomere-1)
 - ⑪ Telomere-2(CCCTTA) ↑ ↑ (Max:5 × Normal Tissue Telomere-2)
- (①-⑤): Omura, Y., 1992-1995; ⑥-⑪: Omura, Y., 1996-1998)

Parameters of Cancer

1. Oncogene C-fos Ab2 : 350ng or higher
2. Integrin $\alpha 5\beta 1$: 350ng or higher
- ✘ Screening & Mapping (60 ng)
3. Acetylcholine Chloride (Normal : $>1500\mu\text{g}$) :
<1pg
4. Hg (Mercury) : 350mg or higher
5. Virus Infection
6. Telomere(TTAGGG)
Cancer Cell Telomere = Normal Cell Telomere $\times$ 5-10
7. 8-OH-dG : 60ng or higher
8. Folic Acid : < 1zg 9. Ghrelin : < 5ng (2ng)
10. IGF-1 : < 5ng (2ng)
11. Strong Resonance with Microscopic Slide
of Specific Cancer Tissue of Specific Internal Organ

BDORTにおけるDHEAの意義

- DHEAの適量は5mgくらい(個人差がある)
- 1日に1回服用(※中国パセリとはキャンセルする)
- 正常なTelomereを増やす。(500ngくらいまで)
- ThromboxaneB2を減らす
- 癌のマーカー(Integrin $\alpha_5\beta_1$ 等を減少させる)

- 他にもp53(Ab-5)
- Rb(Ab-8)
- CA15-3
- CA125
- TGF- α
- TGF- β
- Glucose
- HVA
- VMA
- asialo GM2 etc.

癌のスクリーニング

Fig. 2B: Schematic diagram of magnified cells through projection light microscope to be examined by indirect Bi-Digital O-Ring Test.

3) Electron Microscopic Indirect Bi-Digital O-Ring Test:

Instead of the photon beam in a light microscope, using an electron beam passing through the close vicinity of specific molecules of the specimen in an electron microscope, the molecular information is transmitted to a magnified fluorescent screen, and the indirect Bi-Digital O-Ring Test was performed through a projected penlight-type soft laser beam with wave length of 670 nm on the magnified intracellular fine structure through an observation glass window. Using magnified fine structure of the cells, by an electron microscope, in various cancer cells of both humans and animals, Oncogen C-fos (AB2) and mercury were found inside of the nucleus. The Hitachi Electron Microscope H7000 was used in this study.

Fig. 3A

Fig. 3B

Fig. 3A: Please note the electron microscope set-up. In the observation window of the electron microscope, the laser beam is projected on the magnified screen of the fine structure of the cells from a hand-held solid-state laser beam generator. In the same hand, the pure reference control substance

Acupuncture &
Electro-Therapeutics
Res. Int. J. Vol.17,
pp.29-46, 1992 (p.33)
より引用

$\alpha 5\beta 1$ Integrin

- Integrins are a large family of important proteins that enable cells to bind and respond to extracellular compounds. They are involved in various cell functions including wound healing, cell differentiation, growth of tumor cells and natural cell death. Jerini has focused on a specific integrin, $\alpha 5\beta 1$, which has been demonstrated in preclinical models to play an essential role in the formation of new blood vessels (angiogenesis). Angiogenesis is a major factor in cancer, rheumatoid arthritis, diabetic nephropathy, psoriasis and a number of other diseases including age-related macular degeneration (AMD). In 2003, Jerini initiated the $\alpha 5\beta 1$ integrin project with two programs: 1) the identification and development of a small molecule suitable for slow release delivery in the treatment of AMD, and 2) the identification of an orally-available small molecule drug candidate for oncology.

»Surface representation of a X-ray structure of $\alpha 5\beta 1$ integrin in complex with a peptidic ligand (pdb115g).

Acetylcholine (ACh) synthesis:

FIG. 5. Schematic of the telomere hypothesis of cell aging and immortalization. Changes in telomere length with passage of time (in cell divisions) are represented for germ line, normal somatic, and transformed cells. Events during early embryonic development at the beginning of the time axis are uncertain (dotted lines). Telomerase is presumably activated at some point in gametogenesis because telomere length is maintained in germ-line cells, but it is not known whether telomerase is active in the early embryo before germ-line development. In contrast, the observed decrease in telomere length and lack of telomerase activity in normal somatic cells suggest repression of telomerase in these cells. At the Hayflick limit (M1), we assume that one or more telomeres have lost a threshold amount of TTAGGG, signaling a checkpoint in cell growth (see text). Mean telomere length at this point is designated T1. Partially transformed cells that bypass this checkpoint without activation of telomerase continue to lose telomeres until "crisis" (M2), when most cells have critically short telomeres on many chromosomes (mean telomere length = T2). Cells that activate telomerase, presumably by mutation, may survive crisis. Telomeres can then be stably maintained at any length. (Graph was modified from ref. 13.)

8-OH-dG

活性酸素による8-ヒドロキングアミン生成と修復

葉酸(Folic Acid)

遺伝子の修復因子

癌で低下する

Ki-67 抗原

Ki-67 抗原は細胞周期の G₁/S/G₂/M 期に発現しており、G₀ 期における発現は認められません。また、増殖細胞（培養細胞）では全細胞周期において Ki-67 抗原の発現が認められます。休止期 (G₀) を除く細胞周期の全周期 (G₁ 期, S 期, G₂ 期, M 期) で発現している核タンパクです。Ki-67 抗原に対する抗体により、正常細胞および腫瘍細胞の細胞増殖活性を直接モニターすることが可能です。正常および**多発性骨髄腫、軟部肉腫、前立腺癌、腺癌、乳癌・メラノーマ・NHL・神経芽細胞腫・腎癌**など各種腫瘍の増殖分画を直接モニターするのに非常に有用

Anti-cyclin E

使用目的: 乳癌、消化器癌、大腸癌、子宮癌、卵巣癌等の良性、悪性の判別及び進行度のチェック

(図) 細胞周期と制御因子
(『肝臓病学 Basic Science』(医学書館刊)より)

◇ サイクリンE濃度が高い乳癌患者は予後が悪い University of Texas

「細胞分裂に関わるタンパク質・サイクリンE、サイクリンD1、サイクリンD3、HER2/Neu濃度と死亡率の相関を乳癌患者395人を対象にして調べたところ、サイクリンEが高濃度の患者は低濃度の患者に比べて癌の診断後11年以内に死亡する確率が13倍高い」という研究結果が、2002年11月14日のNEJM誌に発表されました。試験に参加した患者395人のうちステージ1(初期)の乳癌患者が114人いました。この114人中、サイクリンEが低濃度だった患者102人は診断後5年以内に1人も死亡しませんでした。一方サイクリンEが高濃度であった患者12人は診断後5年以内に全員亡くなりました。この結果から、サイクリンE濃度が高い乳癌患者はより強力な治療をすべきであると考えられます。Cyclin E and Survival in Patients with Breast Cancer
NEJM Volume 347:1566-1575 November 14, 2002 Number 20

VEGF · · · Marker of Angiogenesis

The VEGF binds to Flk-1/KDR (tyrosine kinase) receptors on the endothelial cell membranes.

cutaway illustration of an endothelial cell

IGF-1

前立腺がん、乳がん、肺がん、大腸がんのような、何種類かの癌リスクを高める

<5ng

癌の パラメーター

各種癌のパラメーター

各種癌のパラメーター

癌のパラメーター

- P53(Ab-5)
- Integrin $\alpha 5 \beta 1$
- Oncogene C-fos Ab2
- Acetylcholine
- Hg
- Ki-67
- Dopamine

DHEA

- EPA/DHA (EPA180mg + DHA120mg)
- 100mg of Cilantro Tablet
- 100 μ g of Folic Acid

- 75-100mg of Acetyl-L-Carnitine
- 50mg of Lipoic Acid

July 18, 2001

Integrindsβ₁

○: 60ng
⊙: 200ng
⊙: 300ng

Integrindsβ₁: 60ng
Oncogen C-fos Ab: 60ng

Hg: 150ng
Ki67: 100ng
ACh: 50pg

Hepatitis C Monoclonal Antibody: 50ng

ACh 10Zg
1. Integrindsβ₁ 60ng

at ⊙ 5Zg 300ng

2. Oncogen C-fos Ab 2300ng

3. Hg: 200ng

4. Acalylcholin: 10Zg (10⁻²⁰g)

5. Ki67: 100pg (10⁻²⁰g) 200pg

6. P53 Ab 5: 100ng 10ng

7. Virus: 10ng

8. Dopamina 50pg 5μg (N: 1~3pg)

Hepatoma:

チロシン

カテコールアミンの生合成・代謝経路(褐色細胞腫)

• 癌の抗血管新生治療におけるドーパミンの役割

01/07/02 原文の日付は2001年5月31日

- パーキンソン病治療に用いられる神経伝達物質ドーパミンは抗癌治療においても有効か
- 著者: Anthea Hammond博士
- 「Nature Medicine」5月号において、すでにパーキンソン病の治療に臨床使用されている神経伝達物質ドーパミンが、抗血管新生作用をもつことが明らかにされた。これにより、新たに癌や慢性関節リウマチの治療にも、ドーパミンを適用できることが示唆される。パーキンソン病および精神分裂病両方の病因に関係する神経伝達物質であるドーパミンの作用により、数種の悪性腫瘍の成長が抑制されることが、マウスにおいて明らかにされた。ドーパミンD2受容体が内皮細胞上に存在するため、Basuらは、ドーパミンの抗癌作用は、その抗血管新生活性の結果生じているのだらうと論じた。癌の血管新生においてVEGFが重要な役割を果たすことから、Basuらは、ドーパミンがVEGFにより誘導された血管新生を抑制するかについて検討した。

6. 癌におけるAcetylcholineの濃度

July 18, 2001

Integrinds β

- o: 60mg
- ⊙: 200mg
- ⊙: 300mg

Integrinds β : 60mg
Oncagen C-her β : 60mg

Hg: 150mg
Ki67: 100mg

ACh: 50 μ g

Hepatitis C Monoclonal Antibody: 50mg

ACH 102g
52g

1. Integrinds β 60mg
300mg

ACH 102g
52g

ACH 102g
52g

- 2. Oncagen C-her β Ab 2300
- 3. Hg: 200mg
- 4. Acatylcholin: 102g
- 5. Ki67: 100mg
- 6. P53Ab 5: 100mg
- 7. Virus: HHHH
- 8. Dopamina 50 μ g 5 μ g (N: 1-3g)

Hepatoma:

カテコールアミンの代謝経路(神経芽細胞腫)

- VMA/HVA比が低いものはβ-水酸化酵素を欠き、より未分化で予後が悪い

The ErbB Family and Ligands

Activated EGFR-TK

A Pivotal Driver of Malignancy ¹⁻⁶

Adapted with permission from Baselga J. *Signal*. 2000;1:12-21. 1. Raymond E et al. *Drugs*. 2000;60(suppl 1):15-23. 2. Woodburn JR. *Pharmacol Ther*. 1999;82:241-250. 3. Wells A. *Int J Biochem Cell Biol*. 1999;31:637-643. 4. Hanahan D, Weinberg RA. *Cell*. 2000;100:70. 5. Balaban N et al. *Biochim Biophys Acta*. 1996;1314:147-156. 6. Akimoto T et al. *Clin Cancer Res*. 1999;5:2884-2890.

EGFR Expression in Selected Human Tumors

Colon	25-77%
Head and Neck	95-100%
Pancreatic	30-89%
NSCLC	40-80%
Renal cell carcinoma	50-90%
Breast	14-91% (45%)
Ovarian	35-70%
Glioma	40-63%
Bladder	31-48%

Tumours showing high EGFR expression

- NSCLC 40-80%
- Prostate 40-80%
- Gastric 33-74%
- Breast 14-91%
- Head & neck 90-100%
- Colorectal 25-77%
- Pancreatic 30-50%
- Ovarian 35-70%

EGFR expression in human tumours

High expression is generally associated with

- Invasion
- Metastasis
- Late-stage disease
- Chemotherapy resistance
- Hormonal therapy resistance
- Poor outcome